

Observation of the Doubly Charmed Baryon Ξ_{cc}^{++}

LHCb Collaboration

DOI:

[10.1103/PhysRevLett.119.112001](https://doi.org/10.1103/PhysRevLett.119.112001)

License:

Creative Commons: Attribution (CC BY)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

LHCb Collaboration 2017, 'Observation of the Doubly Charmed Baryon Ξ_{cc}^{++} ', *Physical Review Letters*, vol. 119, no. 11, 112001. <https://doi.org/10.1103/PhysRevLett.119.112001>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

checked on 15/1/19

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.



Observation of the Doubly Charmed Baryon Ξ_{cc}^{++}

R. Aaij *et al.**

(LHCb Collaboration)

(Received 6 July 2017; revised manuscript received 2 August 2017; published 11 September 2017)

A highly significant structure is observed in the $\Lambda_c^+ K^- \pi^+ \pi^+$ mass spectrum, where the Λ_c^+ baryon is reconstructed in the decay mode $p K^- \pi^+$. The structure is consistent with originating from a weakly decaying particle, identified as the doubly charmed baryon Ξ_{cc}^{++} . The difference between the masses of the Ξ_{cc}^{++} and Λ_c^+ states is measured to be $1334.94 \pm 0.72(\text{stat.}) \pm 0.27(\text{syst.}) \text{ MeV}/c^2$, and the Ξ_{cc}^{++} mass is then determined to be $3621.40 \pm 0.72(\text{stat.}) \pm 0.27(\text{syst.}) \pm 0.14(\Lambda_c^+) \text{ MeV}/c^2$, where the last uncertainty is due to the limited knowledge of the Λ_c^+ mass. The state is observed in a sample of proton-proton collision data collected by the LHCb experiment at a center-of-mass energy of 13 TeV, corresponding to an integrated luminosity of 1.7 fb^{-1} , and confirmed in an additional sample of data collected at 8 TeV.

DOI: 10.1103/PhysRevLett.119.112001

The quark model [1–3] predicts the existence of multiplets of baryon and meson states. Those states composed of the lightest four quarks (u , d , s , c) form SU(4) multiplets [4]. Numerous states with charm quantum number $C = 0$ or $C = 1$ have been discovered, including all of the expected $q\bar{q}$ and qqq ground states [5]. Three weakly decaying qqq states with $C = 2$ are expected: one isospin doublet ($\Xi_{cc}^{++} = ccu$ and $\Xi_{cc}^{+} = ccd$) and one isospin singlet ($\Omega_{cc}^{+} = ccs$), each with spin parity $J^P = 1/2^+$. The properties of these baryons have been calculated with a variety of theoretical models. In most cases, the masses of the Ξ_{cc} states are predicted to lie in the range 3500 to 3700 MeV/c^2 [6–33]. The masses of the Ξ_{cc}^{++} and Ξ_{cc}^{+} states are expected to differ by only a few MeV/c^2 , due to approximate isospin symmetry [34–36]. Most predictions for the lifetime of the Ξ_{cc}^{++} baryon are in the range 50 to 250 fs, and the lifetime of the Ξ_{cc}^{+} baryon is expected to be three to four times longer at 200 to 700 fs [10,11,19,24,37–40]. While both are expected to be produced at hadron colliders [41–43], the longer lifetime of the Ξ_{cc}^{++} baryon should make it significantly easier to observe than the Ξ_{cc}^{+} baryon in such experiments, due to the use of real-time (online) event-selection requirements designed to reject backgrounds originating from the primary interaction point.

Experimentally, there is a long-standing puzzle in the Ξ_{cc} system. Observations of the Ξ_{cc}^{+} baryon at a mass of $3519 \pm 2 \text{ MeV}/c^2$ with signal yields of 15.9 events over 6.1 ± 0.5 background in the final state $\Lambda_c^+ K^- \pi^+$ (6.3σ significance),

and 5.62 events over 1.38 ± 0.13 background in the final state $p D^+ K^-$ (4.8σ significance) were reported by the SELEX Collaboration [44,45]. Their results included a number of unexpected features, notably a short lifetime and a large production rate relative to that of the singly charmed Λ_c^+ baryon. The lifetime was stated to be shorter than 33 fs at the 90% confidence level, and SELEX concluded that 20% of all Λ_c^+ baryons observed by the experiment originated from Ξ_{cc}^{+} decays, implying a relative Ξ_{cc} production rate several orders of magnitude larger than theoretical expectations [11]. Searches from the FOCUS [46], BABAR [47], and Belle [48] experiments did not find evidence for a state with the properties reported by SELEX, and neither did a search at LHCb with data collected in 2011 corresponding to an integrated luminosity of 0.65 fb^{-1} [49]. However, because the production environments at these experiments differ from that of SELEX, which studied collisions of a hyperon beam on fixed nuclear targets, these null results do not exclude the original observations.

This Letter presents the observation of the Ξ_{cc}^{++} baryon [50] via the decay mode $\Lambda_c^+ K^- \pi^+ \pi^+$ (Fig. 1), which is expected to have a branching fraction of up to 10% [51]. The Λ_c^+ baryon is reconstructed in the final state $p K^- \pi^+$.




FIG. 1. Example Feynman diagram contributing to the decay $\Xi_{cc}^{++} \rightarrow \Lambda_c^+ K^- \pi^+ \pi^+$.

*Full author list given at the end of the article.

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International license](#). Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI.

The data consist of pp collisions collected by the LHCb experiment at the Large Hadron Collider at CERN with a center-of-mass energy of 13 TeV taken in 2016, corresponding to an integrated luminosity of 1.7 fb^{-1} .

The LHCb detector is a single-arm forward spectrometer covering the pseudorapidity range $2 < \eta < 5$, designed for the study of particles containing b or c quarks, and is described in detail in Refs. [52,53]. The detector elements most relevant to this analysis are a silicon-strip vertex detector surrounding the pp interaction region, a tracking system that provides a measurement of the momentum of charged particles, and two ring-imaging Cherenkov detectors [54] that are able to discriminate between different species of charged hadrons. The on-line event selection is performed by a trigger that consists of a hardware stage, which is based on information from the calorimeter and muon systems, followed by a software stage, which fully reconstructs the event [55]. The on-line reconstruction incorporates near-real-time alignment and calibration of the detector [56], which in turn allows the reconstruction of the Ξ_{cc}^{++} decay to be performed entirely in the trigger software.

The reconstruction of $\Xi_{cc}^{++} \rightarrow \Lambda_c^+ K^- \pi^+ \pi^+$ decays proceeds as follows. Candidate $\Lambda_c^+ \rightarrow p K^- \pi^+$ decays are reconstructed from three charged particles that form a good-quality vertex and that are inconsistent with originating from any pp collision primary vertex (PV). The PV of any single particle is defined to be the PV with respect to which the particle has the smallest impact parameter χ^2 (χ_{IP}^2), which is the difference in χ^2 of the PV fit with and without the particle in question. The Λ_c^+ vertex is required to be displaced from its PV by a distance corresponding to a proper decay time greater than 150 fs. The Λ_c^+ candidate is then combined with three additional charged particles to form a $\Xi_{cc}^{++} \rightarrow \Lambda_c^+ K^- \pi^+ \pi^+$ candidate. These additional particles must form a good-quality vertex with the Λ_c^+ candidate, and the Λ_c^+ decay vertex must be downstream of the Ξ_{cc}^{++} vertex. Each of the six final-state particles is required to pass track-quality requirements, to have hadron-identification information consistent with the appropriate hypothesis (p , K , or π), and to have transverse momentum $p_T > 500 \text{ MeV}/c$. To avoid duplicate tracks, the angle between each pair of final-state particles with the same charge is required to be larger than 0.5 mrad . The Ξ_{cc}^{++} candidate must have $p_T > 4 \text{ GeV}/c$ and must be consistent with originating from its PV. The selection above includes criteria applied in the trigger software, plus additional requirements chosen based on simulated signal events and a control sample of data. Simulated signal events are produced with the standard LHCb simulation software [57–63] interfaced to a dedicated generator, GENXICC [64–66], for Ξ_{cc}^{++} baryon production. In the simulation, the Ξ_{cc}^{++} mass and lifetime are assumed to be $3600 \text{ MeV}/c^2$ and 333 fs . The background control sample consists of wrong-sign (WS) $\Lambda_c^+ K^- \pi^+ \pi^-$ combinations.

The background level is further reduced with a multivariate selector based on the multilayer perceptron algorithm [67]. The selector is trained with simulated signal events and with the WS control sample of data to represent the background. For both signal and background training samples, candidates are required to pass the selection described above and to fall within a signal search region defined as $2270 < m_{\text{cand}}(\Lambda_c^+) < 2306 \text{ MeV}/c^2$ and $3300 < m_{\text{cand}}(\Xi_{cc}^{++}) < 3800 \text{ MeV}/c^2$, where $m_{\text{cand}}(\Lambda_c^+)$ is the reconstructed mass of the Λ_c^+ candidate, $m_{\text{cand}}(\Xi_{cc}^{++}) \equiv m(\Lambda_c^+ K^- \pi^+ \pi^+) - m_{\text{cand}}(\Lambda_c^+) + m_{\text{PDG}}(\Lambda_c^+)$, $m(\Lambda_c^+ K^- \pi^+ \pi^+)$ is the reconstructed mass of the $\Lambda_c^+ K^- \pi^+ \pi^+$ combination, and $m_{\text{PDG}}(\Lambda_c^+) = 2286.46 \pm 0.14 \text{ MeV}/c^2$ is the known value of the Λ_c^+ mass [5]. The $m_{\text{cand}}(\Lambda_c^+)$ window corresponds to approximately ± 3 times the Λ_c^+ mass resolution. The use of $m_{\text{cand}}(\Xi_{cc}^{++})$ rather than $m(\Lambda_c^+ K^- \pi^+ \pi^+)$ cancels fluctuations in the reconstructed Λ_c^+ mass to first order, and thereby improves the Ξ_{cc}^{++} mass resolution by approximately 40%.

Based on studies with simulated events and control samples of data, ten input variables that together provide good discrimination between signal and background candidates are used in the multivariate selector. They are as follows: the χ^2 per degree of freedom of each of the Λ_c^+ vertex fit, the Ξ_{cc}^{++} vertex fit, and a kinematic refit [68] of the Ξ_{cc}^{++} decay chain requiring it to originate from its PV; the smallest p_T of the three decay products of the Λ_c^+ ; the smallest p_T of the four decay products of the Ξ_{cc}^{++} ; the scalar sum of the p_T of the four decay products of the Ξ_{cc}^{++} ; the angle between the Ξ_{cc}^{++} momentum vector and the direction from the PV to the Ξ_{cc}^{++} decay vertex; the flight distance χ^2 between the PV and the Ξ_{cc}^{++} decay vertex; the χ_{IP}^2 of the Ξ_{cc}^{++} with respect to its PV; and the smallest χ_{IP}^2 of the decay products of the Ξ_{cc}^{++} with respect to its PV. Here, the flight distance χ^2 is defined as the χ^2 of the hypothesis that the Ξ_{cc}^{++} decay vertex coincides with its PV. Candidates are retained for analysis only if their multivariate selector output values exceed a threshold chosen by maximizing the expected value of the figure of merit $\varepsilon/(\frac{5}{2} + \sqrt{B})$ [69], where ε is the estimated signal efficiency and B is the estimated number of background candidates underneath the signal peak. The quantity B is computed with the WS control sample and, purely for the purposes of this optimization, it is calculated in a window centered at a mass of $3600 \text{ MeV}/c^2$ and of half-width $12.5 \text{ MeV}/c^2$ (corresponding to approximately twice the expected resolution). Its evaluation takes into account the difference in background rates between the $\Lambda_c^+ K^- \pi^+ \pi^+$ signal mode and the WS sample, scaling the WS background by the ratio seen in data in the sideband regions $3200 < m_{\text{cand}}(\Xi_{cc}^{++}) < 3300 \text{ MeV}/c^2$ and $3800 < m_{\text{cand}}(\Xi_{cc}^{++}) < 3900 \text{ MeV}/c^2$. The performance of the multivariate selector is also tested for simulated signal events under other lifetime hypotheses; while the signal efficiency increases with the lifetime, it is




FIG. 2. Mass spectra of (upper) Λ_c^+ and (lower) Ξ_{cc}^{++} candidates. The full selection is applied, except for the Λ_c^+ mass requirement in the case of the upper plot. For the Λ_c^+ mass distribution the (cross-hatched) signal and (vertical line) sideband regions are indicated; to avoid duplication, the histogram is filled only once in events that contain more than one Ξ_{cc}^{++} candidate. In the lower plot the right-sign (RS) signal sample $\Xi_{cc}^{++} \rightarrow \Lambda_c^+ K^- \pi^+ \pi^+$ is shown, along with the control samples: Λ_c^+ sideband (SB) $\Lambda_c^+ K^- \pi^+ \pi^+$ candidates and wrong-sign (WS) $\Lambda_c^+ K^- \pi^+ \pi^-$ candidates, normalized to have the same area as the RS sample in the $m_{\text{cand}}(\Xi_{cc}^{++})$ sidebands.

found that the training obtained for 333 fs is close to optimal (i.e., gives comparable performance to a training optimized for the new lifetime hypothesis) even for much shorter or longer lifetimes.

After the multivariate selection is applied, events may still contain more than one Ξ_{cc}^{++} candidate in the signal search region. Based on studies of simulation and the control data sample, no peaking background arises due to multiple candidates except for the special case in which the candidates are formed from the same six decay products but two of the decay products are interchanged (e.g., the K^- particle from the Ξ_{cc}^{++} decay and the K^- particle from the Λ_c^+ decay). In such instances, one of the candidates is chosen at random to be retained and all others are discarded. In the remaining events, the fraction




FIG. 3. Invariant mass distribution of $\Lambda_c^+ K^- \pi^+ \pi^+$ candidates with fit projections overlaid.

that has more than one Ξ_{cc}^{++} candidate in the range 3300–3800 MeV/ c^2 is approximately 8%.

The selection described above is then applied to data in the search region. Figure 2 shows the Λ_c^+ mass distribution, and the Ξ_{cc}^{++} mass spectra for candidates in the mass range $2270 < m_{\text{cand}}(\Lambda_c^+) < 2306$ MeV/ c^2 . A structure is visible in the signal mode at a mass of approximately 3620 MeV/ c^2 . No significant structure is visible in the WS control sample, or for events in the Λ_c^+ mass sidebands. To measure the properties of the structure, an unbinned extended maximum likelihood fit is performed to the invariant mass distribution in the restricted $\Lambda_c^+ K^- \pi^+ \pi^+$ mass window of 3620 ± 150 MeV/ c^2 (Fig. 3). The peaking structure is empirically described by a Gaussian function plus a modified Gaussian function with power-law tails on both sides [70]. All peak parameters are fixed to values obtained from simulation apart from the mass, yield, and an overall resolution parameter. The background is described by a second-order polynomial with parameters free to float in the fit. The signal yield is measured to be 313 ± 33 , corresponding to a local statistical significance in excess of 12σ when evaluated with a likelihood ratio test. The fitted resolution parameter is 6.6 ± 0.8 MeV/ c^2 , consistent with simulation. The same structure is also observed in the $\Lambda_c^+ K^- \pi^+ \pi^+$ spectrum in a pp data sample collected by LHCb at $\sqrt{s} = 8$ TeV (see the Supplemental Material [71] for results from the 8 TeV cross-check sample). The local statistical significance of the peak in the 8 TeV sample is above 7σ , and its mass is consistent with that in the 13 TeV data sample.

Additional cross-checks are performed confirming the robustness of the observation. The significance of the structure in the $\Lambda_c^+ K^- \pi^+ \pi^+$ final state remains above 12σ when fixing the resolution parameter in the invariant mass fit to the value obtained from simulation, changing the threshold value for the multivariate selector, removing events containing multiple candidates in the fitted mass

TABLE I. Systematic uncertainties on the Ξ_{cc}^{++} mass measurement.

Source	Value (MeV/ c^2)
Momentum-scale calibration	0.22
Selection bias correction	0.14
Unknown Ξ_{cc}^{++} lifetime	0.06
Mass fit model	0.07
Sum of above in quadrature	0.27
Λ_c^+ mass uncertainty	0.14

range, or using an alternative selection without a multivariate classifier. The significance also remains above 12σ in a subsample of candidates for which the reconstructed decay time exceeds five times its uncertainty. This is consistent with a weakly decaying state and inconsistent with the strong decay of a resonance. No fake peaking structures are observed in the control samples when requiring various intermediate resonances to be present (ρ^0 , K^{*0} , Σ_c^0 , Σ_c^{++} , Λ_c^{++}) nor are they observed when combining Ξ_{cc}^{++} and Λ_c^+ decay products. The contributions of misidentified $D_s^+ \rightarrow K^+ K^- \pi^+$ and $D^+ \rightarrow K^- \pi^+ \pi^+$ decays are found to be negligible.

The sources of systematic uncertainty affecting the measurement of the Ξ_{cc}^{++} mass (Table I) include the momentum-scale calibration, the event selection, the unknown Ξ_{cc}^{++} lifetime, the invariant mass fit model, and the uncertainty on the Λ_c^+ mass. The momentum scale is calibrated with samples of $J/\psi \rightarrow \mu^+ \mu^-$ and $B^+ \rightarrow J/\psi K^+$ decays [72,73]. After calibration, an uncertainty of $\pm 0.03\%$ is assigned, which corresponds to a systematic uncertainty of $0.22 \text{ MeV}/c^2$ on the reconstructed Ξ_{cc}^{++} mass. The selection procedure is more efficient for vertices that are well separated from the PV, and therefore preferentially retains longer-lived Ξ_{cc}^{++} candidates. Because of a correlation between the reconstructed decay time and the reconstructed mass, this induces a positive bias on the mass for both Ξ_{cc}^{++} and Λ_c^+ candidates. The effect is studied with simulation and the bias on the Ξ_{cc}^{++} mass is determined to be $+0.45 \pm 0.14 \text{ MeV}/c^2$ (assuming a lifetime of 333 fs), where the uncertainty is due to the limited size of the simulation sample. A corresponding correction is applied to the fitted value in data. To validate this procedure, the Λ_c^+ mass in an inclusive sample is measured and corrected in the same way; after the correction, the Λ_c^+ mass is found to agree with the known value [5]. The bias on the Ξ_{cc}^{++} mass depends on the unknown Ξ_{cc}^{++} lifetime, introducing a further source of uncertainty on the correction. This is estimated by repeating the procedure for other Ξ_{cc}^{++} lifetime hypotheses between 200 and 700 fs. The largest deviation in the correction, $0.06 \text{ MeV}/c^2$, is taken as an additional systematic uncertainty. Final-state photon radiation also causes a bias in the measured mass, which is determined to be $-0.05 \text{ MeV}/c^2$ with simulation [61]. The uncertainty on this correction is approximately $0.01 \text{ MeV}/c^2$ and is

neglected. The dependence of the measurement on the fit model is estimated by varying the shape parameters that are fixed according to simulation, by using alternative signal and background models, and by repeating the fits in different mass ranges. The largest deviation seen in the mass, $0.07 \text{ MeV}/c^2$, is assigned as a systematic uncertainty. Finally, since the Ξ_{cc}^{++} mass is measured relative to the Λ_c^+ mass, the uncertainty of $0.14 \text{ MeV}/c^2$ on the world-average value of the latter is included. After taking these systematic effects into account and combining their uncertainties (except that on the Λ_c^+ mass) in quadrature, the Ξ_{cc}^{++} mass is measured to be $3621.40 \pm 0.72(\text{stat.}) \pm 0.27(\text{syst.}) \pm 0.14(\Lambda_c^+) \text{ MeV}/c^2$. The mass difference between the Ξ_{cc}^{++} and Λ_c^+ states is $1334.94 \pm 0.72(\text{stat.}) \pm 0.27(\text{syst.}) \text{ MeV}/c^2$.

In summary, a highly significant structure is observed in the final state $\Lambda_c^+ K^- \pi^+ \pi^+$ in a pp data sample collected by LHCb at $\sqrt{s} = 13 \text{ TeV}$, with a signal yield of 313 ± 33 . The mass of the structure is measured to be $3621.40 \pm 0.72(\text{stat.}) \pm 0.27(\text{syst.}) \pm 0.14(\Lambda_c^+) \text{ MeV}/c^2$, where the last uncertainty is due to the limited knowledge of the Λ_c^+ mass, and its width is consistent with experimental resolution. The structure is confirmed with consistent mass in a data set collected by LHCb at $\sqrt{s} = 8 \text{ TeV}$. The signal candidates have significant decay lengths, and the signal remains highly significant after a minimum lifetime requirement of approximately five times the expected decay-time resolution is imposed. This state is therefore incompatible with a strongly decaying particle but is consistent with the expectations for the weakly decaying Ξ_{cc}^{++} baryon. The mass of the observed Ξ_{cc}^{++} state is greater than that of the Ξ_{cc}^+ peaks reported by the SELEX Collaboration [44,45] by $103 \pm 2 \text{ MeV}/c^2$. This difference would imply an isospin splitting vastly larger than that seen in any other baryon system and is inconsistent with the expected size of a few MeV/c^2 [34–36]. Consequently, while the state reported here is consistent with most theoretical expectations for the Ξ_{cc}^{++} baryon, it is inconsistent with being an isospin partner to the Ξ_{cc}^+ state reported previously by the SELEX Collaboration.

We thank Chao-Hsi Chang, Cai-Dian Lü, Xing-Gang Wu, and Fu-Sheng Yu for frequent and interesting discussions on the production and decays of double-heavy-flavor baryons. We express our gratitude to our colleagues in the CERN accelerator departments for the excellent performance of the LHC. We thank the technical and administrative staff at the LHCb institutes. We acknowledge support from CERN and from the national agencies: CAPES, CNPq, FAPERJ, and FINEP (Brazil); MOST and NSFC (China); CNRS/IN2P3 (France); BMBF, DFG, and MPG (Germany); INFN (Italy); NWO (The Netherlands); MNiSW and NCN (Poland); MEN/IFA (Romania); MinES and FASO (Russia); MinEco (Spain); SNSF and SER (Switzerland); NASU (Ukraine); STFC (United Kingdom);

NSF (USA). We acknowledge the computing resources that are provided by CERN, IN2P3 (France), KIT and DESY (Germany), INFN (Italy), SURF (The Netherlands), PIC (Spain), GridPP (United Kingdom), RRCKI and Yandex LLC (Russia), CSCS (Switzerland), IFIN-HH (Romania), CBPF (Brazil), PL-GRID (Poland) and OSC (USA). We are indebted to the communities behind the multiple open source software packages on which we depend. Individual groups or members have received support from AvH Foundation (Germany), EPLANET, Marie Skłodowska-Curie Actions, and ERC (European Union), Conseil Général de Haute-Savoie, Labex ENIGMASS, and OCEVU, Région Auvergne (France), RFBR and Yandex LLC (Russia), GVA, XuntaGal, and GENCAT (Spain), Herchel Smith Fund, The Royal Society, Royal Commission for the Exhibition of 1851, and the Leverhulme Trust (United Kingdom).

-
- [1] M. Gell-Mann, *Phys. Lett.* **8**, 214 (1964).
 [2] G. Zweig, Report No. CERN-TH-401, 1964.
 [3] G. Zweig in *Developments in the Quark Theory of Hadrons*, edited by D. Lichtenberg and S. Rosen (Hadronic Press, Nonantum, Massachusetts, 1980), Vol. 1, pp. 22–101.
 [4] A. De Rújula, H. Georgi, and S. Glashow, *Phys. Rev. D* **12**, 147 (1975).
 [5] C. Patrignani *et al.* (Particle Data Group), *Chin. Phys. C* **40**, 100001 (2016), and 2017 update.
 [6] S. S. Gershtein, V. V. Kiselev, A. K. Likhoded, and A. I. Onishchenko, *Yad. Fiz.* **63**, 334 (2000) [*Phys. At. Nucl.* **63**, 274 (2000)].
 [7] S. S. Gershtein, V. V. Kiselev, A. K. Likhoded, and A. I. Onishchenko, *Mod. Phys. Lett. A* **14**, 135 (1999).
 [8] C. Itoh, T. Minamikawa, K. Miura, and T. Watanabe, *Phys. Rev. D* **61**, 057502 (2000).
 [9] S. S. Gershtein, V. V. Kiselev, A. K. Likhoded, and A. I. Onishchenko, *Phys. Rev. D* **62**, 054021 (2000).
 [10] K. Anikeev *et al.*, in *Workshop on B physics at the Tevatron: Run II and beyond, Batavia, Illinois, 1999* (2001).
 [11] V. Kiselev and A. Likhoded, *Phys. Usp.* **45**, 455 (2002).
 [12] D. Ebert, R. N. Faustov, V. O. Galkin, and A. P. Martynenko, *Phys. Rev. D* **66**, 014008 (2002).
 [13] D.-H. He, K. Qian, Y.-B. Ding, X.-Q. Li, and P.-N. Shen, *Phys. Rev. D* **70**, 094004 (2004).
 [14] C.-H. Chang, C.-F. Qiao, J.-X. Wang, and X.-G. Wu, *Phys. Rev. D* **73**, 094022 (2006).
 [15] W. Roberts and M. Pervin, *Int. J. Mod. Phys. A* **23**, 2817 (2008).
 [16] A. Valcarce, H. Garcilazo, and J. Vijande, *Eur. Phys. J. A* **37**, 217 (2008).
 [17] J.-R. Zhang and M.-Q. Huang, *Phys. Rev. D* **78**, 094007 (2008).
 [18] Z.-G. Wang, *Eur. Phys. J. A* **45**, 267 (2010).
 [19] M. Karliner and J. L. Rosner, *Phys. Rev. D* **90**, 094007 (2014).
 [20] K.-W. Wei, B. Chen, and X.-H. Guo, *Phys. Rev. D* **92**, 076008 (2015).
 [21] Z.-F. Sun and M. J. Vicente Vacas, *Phys. Rev. D* **93**, 094002 (2016).
 [22] C. Alexandrou and C. Kallidonis, *Phys. Rev. D* **96**, 034511 (2017).
 [23] B. O. Kerbikov, M. I. Polikarpov, and L. V. Shevchenko, *Nucl. Phys.* **B331**, 19 (1990).
 [24] S. Fleck and J.-M. Richard, *Prog. Theor. Phys.* **82**, 760 (1989).
 [25] S. Chernyshev, M. A. Nowak, and I. Zahed, *Phys. Rev. D* **53**, 5176 (1996).
 [26] T. M. Aliev, K. Azizi, and M. Savcı, *Nucl. Phys.* **A895**, 59 (2012).
 [27] Z.-F. Sun, Z.-W. Liu, X. Liu, and S.-L. Zhu, *Phys. Rev. D* **91**, 094030 (2015).
 [28] N. Mathur, R. Lewis, and R. M. Woloshyn, *Phys. Rev. D* **66**, 014502 (2002).
 [29] Y. Namekawa, S. Aoki, K. I. Ishikawa, N. Ishizuka, K. Kanaya, Y. Kuramashi, M. Okawa, Y. Taniguchi, A. Ukawa, N. Ukita, and T. Yoshie (PACS-CS Collaboration), *Phys. Rev. D* **87**, 094512 (2013).
 [30] Z. S. Brown, W. Detmold, S. Meinel, and K. Orginos, *Phys. Rev. D* **90**, 094507 (2014).
 [31] M. Padmanath, R. G. Edwards, N. Mathur, and M. Peardon, *Phys. Rev. D* **91**, 094502 (2015).
 [32] P. Pérez-Rubio, S. Collins, and G. S. Bali, *Phys. Rev. D* **92**, 034504 (2015).
 [33] Y. Liu and I. Zahed, *Phys. Rev. D* **95**, 116012 (2017).
 [34] C.-W. Hwang and C.-H. Chung, *Phys. Rev. D* **78**, 073013 (2008).
 [35] S. J. Brodsky, F.-K. Guo, C. Hanhart, and U.-G. Meißner, *Phys. Lett. B* **698**, 251 (2011).
 [36] M. Karliner and J. L. Rosner, [arXiv:1706.06961](https://arxiv.org/abs/1706.06961).
 [37] B. Guberina, B. Melić, and H. Štefančić, *Eur. Phys. J. C* **9**, 213 (1999).
 [38] V. V. Kiselev, A. K. Likhoded, and A. I. Onishchenko, *Phys. Rev. D* **60**, 014007 (1999).
 [39] C.-H. Chang, T. Li, X.-Q. Li, and Y.-M. Wang, *Commun. Theor. Phys.* **49**, 993 (2008).
 [40] A. V. Berezhnuy and A. K. Likhoded, *Yad. Fiz.* **79**, 151 (2016) [*Phys. At. Nucl.* **79**, 260 (2016)].
 [41] A. V. Berezhnuy, A. K. Likhoded, and M. V. Shevlyagin, *Yad. Fiz.* **58**, 730 (1995) [*Phys. At. Nucl.* **58**, 672 (1995)].
 [42] K. Kolodziej, A. Leike, and R. Ruckl, *Phys. Lett. B* **355**, 337 (1995).
 [43] A. V. Berezhnuy, V. V. Kiselev, A. K. Likhoded, and A. I. Onishchenko, *Phys. Rev. D* **57**, 4385 (1998).
 [44] M. Mattson *et al.* (SELEX Collaboration), *Phys. Rev. Lett.* **89**, 112001 (2002).
 [45] A. Ocherashvili *et al.* (SELEX Collaboration), *Phys. Lett. B* **628**, 18 (2005).
 [46] S. P. Ratti, *Nucl. Phys. B, Proc. Suppl.* **115**, 33 (2003).
 [47] B. Aubert *et al.* (BABAR Collaboration), *Phys. Rev. D* **74**, 011103 (2006).
 [48] R. Chistov *et al.* (Belle Collaboration), *Phys. Rev. Lett.* **97**, 162001 (2006).
 [49] R. Aaij *et al.* (LHCb Collaboration), *J. High Energy Phys.* **12** (2013) 090.
 [50] Inclusion of charge-conjugate processes is implied throughout.
 [51] F.-S. Yu, H.-Y. Jiang, R.-H. Li, C.-D. L. W. Wang, and Z.-X. Zhao, [arXiv:1703.09086](https://arxiv.org/abs/1703.09086).

- [52] A. A. Alves, Jr. *et al.* (LHCb Collaboration), *J. Instrum.* **3**, S08005 (2008).
- [53] R. Aaij *et al.* (LHCb Collaboration), *Int. J. Mod. Phys. A* **30**, 1530022 (2015).
- [54] M. Adinolfi *et al.*, *Eur. Phys. J. C* **73**, 2431 (2013).
- [55] R. Aaij *et al.*, *J. Instrum.* **8**, P04022 (2013).
- [56] G. Dujany and B. Storaci, *J. Phys. Conf. Ser.* **664**, 082010 (2015).
- [57] T. Sjöstrand, S. Mrenna, and P. Skands, *Comput. Phys. Commun.* **178**, 852 (2008).
- [58] T. Sjöstrand, S. Mrenna, and P. Skands, *J. High Energy Phys.* **05** (2006) 026.
- [59] I. Belyaev *et al.*, *J. Phys. Conf. Ser.* **331**, 032047 (2011).
- [60] D. J. Lange, *Nucl. Instrum. Methods Phys. Res., Sect. A* **462**, 152 (2001).
- [61] P. Golonka and Z. Was, *Eur. Phys. J. C* **45**, 97 (2006).
- [62] S. Agostinelli *et al.* (Geant4 Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **506**, 250 (2003); J. Allison, K. Amako, J. Apostolakis, H. Araujo, P. Dubois *et al.* (Geant4 Collaboration), *IEEE Trans. Nucl. Sci.* **53**, 270 (2006).
- [63] M. Clemencic, G. Corti, S. Easo, C. R. Jones, S. Miglioranza, M. Pappagallo, and P. Robbe, *J. Phys. Conf. Ser.* **331**, 032023 (2011).
- [64] C.-H. Chang, J.-X. Wang, and X.-G. Wu, *Comput. Phys. Commun.* **177**, 467 (2007).
- [65] C.-H. Chang, J.-X. Wang, and X.-G. Wu, *Comput. Phys. Commun.* **181**, 1144 (2010).
- [66] X.-Y. Wang and X.-G. Wu, *Comput. Phys. Commun.* **184**, 1070 (2013).
- [67] A. Hoecker, P. Speckmayer, J. Stelzer, J. Therhaag, E. von Toerne, and H. Voss, *Proc. Sci. ACAT* (2007) 040.
- [68] W. D. Hulsbergen, *Nucl. Instrum. Methods Phys. Res., Sect. A* **552**, 566 (2005).
- [69] G. Punzi, in *Statistical Problems in Particle Physics, Astrophysics, and Cosmology*, edited by L. Lyons, R. Mount, and R. Reitmeyer, eConf C030908, 79 (2003).
- [70] T. Skwarnicki, Ph.D. thesis, Institute of Nuclear Physics, Krakow, 1986, DESY-F31-86-02.
- [71] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.119.112001> for results from the 8 TeV cross-check sample.
- [72] R. Aaij *et al.* (LHCb Collaboration), *Phys. Lett. B* **708**, 241 (2012).
- [73] R. Aaij *et al.* (LHCb Collaboration), *J. High Energy Phys.* **06** (2013) 065.

R. Aaij,⁴⁰ B. Adeva,³⁹ M. Adinolfi,⁴⁸ Z. Ajaltouni,⁵ S. Akar,⁵⁹ J. Albrecht,¹⁰ F. Alessio,⁴⁰ M. Alexander,⁵³ A. Alfonso Alberro,³⁸ S. Ali,⁴³ G. Alkhazov,³¹ P. Alvarez Cartelle,⁵⁵ A. A. Alves Jr.,⁵⁹ S. Amato,² S. Amerio,²³ Y. Amhis,⁷ L. An,³ L. Anderlini,¹⁸ G. Andreassi,⁴¹ M. Andreotti,^{17,a} J. E. Andrews,⁶⁰ R. B. Appleby,⁵⁶ F. Archilli,⁴³ P. d'Argent,¹² J. Arnau Romeu,⁶ A. Artamonov,³⁷ M. Artuso,⁶¹ E. Aslanides,⁶ G. Auriemma,²⁶ M. Baalouch,⁵ I. Babuschkin,⁵⁶ S. Bachmann,¹² J. J. Back,⁵⁰ A. Badalov,³⁸ C. Baesso,⁶² S. Baker,⁵⁵ V. Balagura,^{7,b} W. Baldini,¹⁷ A. Baranov,³⁵ R. J. Barlow,⁵⁶ C. Barschel,⁴⁰ S. Barsuk,⁷ W. Barter,⁵⁶ F. Baryshnikov,³² V. Batozskaya,²⁹ V. Battista,⁴¹ A. Bay,⁴¹ L. Beaucourt,⁴ J. Beddow,⁵³ F. Bedeschi,²⁴ I. Bediaga,¹ A. Beiter,⁶¹ L. J. Bel,⁴³ N. Beliy,⁶³ V. Bellee,⁴¹ N. Belloli,^{21,c} K. Belous,³⁷ I. Belyaev,³² E. Ben-Haim,⁸ G. Bencivenni,¹⁹ S. Benson,⁴³ S. Beranek,⁹ A. Berezhnoy,³³ R. Bernet,⁴² D. Berninghoff,¹² E. Bertholet,⁸ A. Bertolin,²³ C. Betancourt,⁴² F. Betti,¹⁵ M.-O. Bettler,⁴⁰ M. van Beuzekom,⁴³ I. Bezshyiko,⁴² S. Bifani,⁴⁷ P. Billoir,⁸ A. Birnkraut,¹⁰ A. Bitadze,⁵⁶ A. Bizzeti,^{18,d} M. B. Bjoern,⁵⁷ T. Blake,⁵⁰ F. Blanc,⁴¹ J. Blouw,^{11,f} S. Blusk,⁶¹ V. Bocci,²⁶ T. Boettcher,⁵⁸ A. Bondar,^{36,e} N. Bondar,³¹ W. Bonivento,¹⁶ I. Bordyuzhin,³² A. Borgheresi,^{21,c} S. Borghi,⁵⁶ M. Borisyak,³⁵ M. Borsato,³⁹ M. Borysova,⁴⁶ F. Bossu,⁷ M. Boubdir,⁹ T. J. V. Bowcock,⁵⁴ E. Bowen,⁴² C. Bozzi,^{17,40} S. Braun,¹² T. Britton,⁶¹ J. Brodzicka,²⁷ D. Brundu,¹⁶ E. Buchanan,⁴⁸ C. Burr,⁵⁶ A. Bursche,^{16,f} J. Buytaert,⁴⁰ W. Byczynski,⁴⁰ S. Cadetdu,¹⁶ H. Cai,⁶⁴ R. Calabrese,^{17,a} R. Calladine,⁴⁷ M. Calvi,^{21,c} M. Calvo Gomez,^{38,g} A. Camboni,³⁸ P. Campana,¹⁹ D. H. Campora Perez,⁴⁰ L. Capriotti,⁵⁶ A. Carbone,^{15,h} G. Carboni,^{25,i} R. Cardinale,^{20,j} A. Cardini,¹⁶ P. Carniti,^{21,c} L. Carson,⁵² K. Carvalho Akiba,² G. Casse,⁵⁴ L. Cassina,^{21,c} L. Castillo Garcia,⁴¹ M. Cattaneo,⁴⁰ G. Cavallero,^{20,40,j} R. Cenci,^{24,k} D. Chamont,⁷ M. Charles,⁸ Ph. Charpentier,⁴⁰ G. Chatzikonstantinidis,⁴⁷ M. Chefdeville,⁴ S. Chen,⁵⁶ S. F. Cheung,⁵⁷ S.-G. Chitic,⁴⁰ V. Chobanova,³⁹ M. Chrzasczcz,^{42,27} A. Chubykin,³¹ P. Ciambrone,¹⁹ X. Cid Vidal,³⁹ G. Ciezarek,⁴³ P. E. L. Clarke,⁵² M. Clemencic,⁴⁰ H. V. Cliff,⁴⁹ J. Closier,⁴⁰ J. Cogan,⁶ E. Cogneras,⁵ V. Cogoni,^{16,f} L. Cojocariu,³⁰ P. Collins,⁴⁰ T. Colombo,⁴⁰ A. Comerma-Montells,¹² A. Contu,⁴⁰ A. Cook,⁴⁸ G. Coombs,⁴⁰ S. Coquereau,³⁸ G. Corti,⁴⁰ M. Corvo,^{17,a} C. M. Costa Sobral,⁵⁰ B. Couturier,⁴⁰ G. A. Cowan,⁵² D. C. Craik,⁵⁸ A. Crocombe,⁵⁰ M. Cruz Torres,⁶² R. Currie,⁵² C. D'Ambrosio,⁴⁰ F. Da Cunha Marinho,² E. Dall'Occo,⁴³ J. Dalseno,⁴⁸ A. Davis,³ O. De Aguiar Francisco,⁵⁴ S. De Capua,⁵⁶ M. De Cian,¹² J. M. De Miranda,¹ L. De Paula,² M. De Serio,^{14,l} P. De Simone,¹⁹ C. T. Dean,⁵³ D. Decamp,⁴ L. Del Buono,⁸ H.-P. Dembinski,¹¹ M. Demmer,¹⁰ A. Dendek,²⁸ D. Derkach,³⁵ O. Deschamps,⁵ F. Dettori,⁵⁴ B. Dey,⁶⁵ A. Di Canto,⁴⁰ P. Di Nezza,¹⁹ H. Dijkstra,⁴⁰ F. Dordei,⁴¹ M. Dorigo,⁴¹ A. Dosil Suárez,³⁹ L. Douglas,⁵³ A. Dovbnya,⁴⁵ K. Dreimanis,⁵⁴ L. Dufour,⁴³ G. Dujany,⁸ P. Durante,⁴⁰ R. Dzhelezadine,³⁷ M. Dziewiecki,¹² A. Dziurda,⁴⁰ A. Dzyuba,³¹ S. Easo,⁵¹ M. Ebert,⁵² U. Egede,⁵⁵ V. Egorychev,³² S. Eidelman,^{36,e}

S. Eisenhardt,⁵² U. Eitschberger,¹⁰ R. Ekelhof,¹⁰ L. Eklund,⁵³ S. Ely,⁶¹ S. Esen,¹² H. M. Evans,⁴⁹ T. Evans,⁵⁷ A. Falabella,¹⁵ N. Farley,⁴⁷ S. Farry,⁵⁴ R. Fay,⁵⁴ D. Fazzini,^{21,c} L. Federici,²⁵ D. Ferguson,⁵² G. Fernandez,³⁸ P. Fernandez Declara,⁴⁰ A. Fernandez Prieto,³⁹ F. Ferrari,¹⁵ F. Ferreira Rodrigues,² M. Ferro-Luzzi,⁴⁰ S. Filippov,³⁴ R. A. Fini,¹⁴ M. Fiore,^{17,a} M. Fiorini,^{17,a} M. Firlej,²⁸ C. Fitzpatrick,⁴¹ T. Fiutowski,²⁸ F. Fleuret,^{7,m} K. Fohl,⁴⁰ M. Fontana,^{16,40} F. Fontanelli,^{20,j} D. C. Forshaw,⁶¹ R. Forty,⁴⁰ V. Franco Lima,⁵⁴ M. Frank,⁴⁰ C. Frei,⁴⁰ J. Fu,^{22,n} W. Funk,⁴⁰ E. Furfaro,^{25,i} C. Färber,⁴⁰ E. Gabriel,⁵² A. Gallas Torreira,³⁹ D. Galli,^{15,h} S. Gallorini,²³ S. Gambetta,⁵² M. Gandelman,² P. Gandini,⁵⁷ Y. Gao,³ L. M. Garcia Martin,⁷⁰ J. García Pardiñas,³⁹ J. Garra Tico,⁴⁹ L. Garrido,³⁸ P. J. Garsed,⁴⁹ D. Gascon,³⁸ C. Gaspar,⁴⁰ L. Gavardi,¹⁰ G. Gazzoni,⁵ D. Gerick,¹² E. Gersabeck,¹² M. Gersabeck,⁵⁶ T. Gershon,⁵⁰ Ph. Ghez,⁴ S. Giani,⁴¹ V. Gibson,⁴⁹ O. G. Girard,⁴¹ L. Giubega,³⁰ K. Gizdov,⁵² V. V. Gligorov,⁸ D. Golubkov,³² A. Golutvin,^{55,40} A. Gomes,^{1,o} I. V. Gorelov,³³ C. Gotti,^{21,c} E. Govorkova,⁴³ J. P. Grabowski,¹² R. Graciani Diaz,³⁸ L. A. Granado Cardoso,⁴⁰ E. Graugés,³⁸ E. Graverini,⁴² G. Graziani,¹⁸ A. Grecu,³⁰ R. Greim,⁹ P. Griffith,¹⁶ L. Grillo,^{21,40,c} L. Gruber,⁴⁰ B. R. Gruberg Cazon,⁵⁷ O. Grünberg,⁶⁷ E. Gushchin,³⁴ Yu. Guz,³⁷ T. Gys,⁴⁰ C. Göbel,⁶² T. Hadavizadeh,⁵⁷ C. Hadjivasiliou,⁵ G. Haefeli,⁴¹ C. Haen,⁴⁰ S. C. Haines,⁴⁹ B. Hamilton,⁶⁰ X. Han,¹² T. Hancock,⁵⁷ S. Hansmann-Menzemer,¹² N. Harnew,⁵⁷ S. T. Harnew,⁴⁸ J. Harrison,⁵⁶ C. Hasse,⁴⁰ M. Hatch,⁴⁰ J. He,⁶³ M. Hecker,⁵⁵ K. Heinicke,¹⁰ A. Heister,⁹ K. Hennessy,⁵⁴ P. Henrard,⁵ L. Henry,⁷⁰ E. van Herwijnen,⁴⁰ M. Heß,⁶⁷ A. Hicheur,² D. Hill,⁵⁷ C. Hombach,⁵⁶ P. H. Hopchev,⁴¹ Z.-C. Huard,⁵⁹ W. Hulsbergen,⁴³ T. Humair,⁵⁵ M. Hushchyn,³⁵ D. Hutchcroft,⁵⁴ P. Ibis,¹⁰ M. Idzik,²⁸ P. Ilten,⁵⁸ R. Jacobsson,⁴⁰ J. Jalocha,⁵⁷ E. Jans,⁴³ A. Jawahery,⁶⁰ F. Jiang,³ M. John,⁵⁷ D. Johnson,⁴⁰ C. R. Jones,⁴⁹ C. Joram,⁴⁰ B. Jost,⁴⁰ N. Jurik,⁵⁷ S. Kandybei,⁴⁵ M. Karacson,⁴⁰ J. M. Kariuki,⁴⁸ S. Karodia,⁵³ N. Kazeev,³⁵ M. Kecke,¹² M. Kelsey,⁶¹ M. Kenzie,⁴⁹ T. Ketel,⁴⁴ E. Khairullin,³⁵ B. Khanji,¹² C. Khurewathanakul,⁴¹ T. Kim,⁹ S. Klaver,⁵⁶ K. Klimaszewski,²⁹ T. Klimkovich,¹¹ S. Koliev,⁴⁶ M. Kolpin,¹² I. Komarov,⁴¹ R. Kopečna,¹² P. Koppenburg,⁴³ A. Kosmyntseva,³² S. Kotriakhova,³¹ M. Kozeiha,⁵ M. Kreps,⁵⁰ P. Krokovny,^{36,e} F. Kruse,¹⁰ W. Krzemien,²⁹ W. Kucewicz,^{27,p} M. Kucharczyk,²⁷ V. Kudryavtsev,^{36,e} A. K. Kuonen,⁴¹ K. Kurek,²⁹ T. Kvaratskheliya,^{32,40} D. Lacarrere,⁴⁰ G. Lafferty,⁵⁶ A. Lai,¹⁶ G. Lanfranchi,¹⁹ C. Langenbruch,⁹ T. Latham,⁵⁰ C. Lazzeroni,⁴⁷ R. Le Gac,⁶ J. van Leerdam,⁴³ A. Leflat,^{33,40} J. Lefrançois,⁷ R. Lefèvre,⁵ F. Lemaitre,⁴⁰ E. Lemos Cid,³⁹ O. Leroy,⁶ T. Lesiak,²⁷ B. Leverington,¹² P.-R. Li,⁶³ T. Li,³ Y. Li,⁷ Z. Li,⁶¹ T. Likhomanenko,⁶⁸ R. Lindner,⁴⁰ F. Lionetto,⁴² V. Lisovskyi,⁷ X. Liu,³ D. Loh,⁵⁰ A. Loi,¹⁶ I. Longstaff,⁵³ J. H. Lopes,² D. Lucchesi,^{23,q} M. Lucio Martinez,³⁹ H. Luo,⁵² A. Lupato,²³ E. Luppi,^{17,a} O. Lupton,⁴⁰ A. Lusiani,²⁴ X. Lyu,⁶³ F. Machefert,⁷ F. Maciuc,³⁰ V. Macko,⁴¹ P. Mackowiak,¹⁰ B. Maddock,⁵⁹ S. Maddrell-Mander,⁴⁸ O. Maev,³¹ K. Maguire,⁵⁶ D. Maisuzenko,³¹ M. W. Majewski,²⁸ S. Malde,⁵⁷ A. Malinin,⁶⁸ T. Maltsev,³⁶ G. Manca,^{16,f} G. Mancinelli,⁶ P. Manning,⁶¹ D. Marangotto,^{22,n} J. Maratas,^{5,r} J. F. Marchand,⁴ U. Marconi,¹⁵ C. Marin Benito,³⁸ M. Marinangeli,⁴¹ P. Marino,⁴¹ J. Marks,¹² G. Martellotti,²⁶ M. Martin,⁶ M. Martinelli,⁴¹ D. Martinez Santos,³⁹ F. Martinez Vidal,⁷⁰ D. Martins Tostes,² L. M. Massacrier,⁷ A. Massafferri,¹ R. Matev,⁴⁰ A. Mathad,⁵⁰ Z. Mathe,⁴⁰ C. Matteuzzi,²¹ A. Mauri,⁴² E. Maurice,^{7,m} B. Maurin,⁴¹ A. Mazurov,⁴⁷ M. McCann,^{55,40} A. McNab,⁵⁶ R. McNulty,¹³ J. V. Mead,⁵⁴ B. Meadows,⁵⁹ C. Meaux,⁶ F. Meier,¹⁰ N. Meinert,⁶⁷ D. Melnychuk,²⁹ M. Merk,⁴³ A. Merli,^{22,40,n} E. Michielin,²³ D. A. Milanes,⁶⁶ E. Millard,⁵⁰ M.-N. Minard,⁴ L. Minzoni,¹⁷ D. S. Mitzel,¹² A. Mogini,⁸ J. Molina Rodriguez,¹ T. Mombacher,¹⁰ I. A. Monroy,⁶⁶ S. Monteil,⁵ M. Morandin,²³ M. J. Morello,^{24,k} O. Morgunova,⁶⁸ J. Moron,²⁸ A. B. Morris,⁵² R. Mountain,⁶¹ F. Muheim,⁵² M. Mulder,⁴³ D. Müller,⁵⁶ J. Müller,¹⁰ K. Müller,⁴² V. Müller,¹⁰ P. Naik,⁴⁸ T. Nakada,⁴¹ R. Nandakumar,⁵¹ A. Nandi,⁵⁷ I. Nasteva,² M. Needham,⁵² N. Neri,^{22,40} S. Neubert,¹² N. Neufeld,⁴⁰ M. Neuner,¹² T. D. Nguyen,⁴¹ C. Nguyen-Mau,^{41,s} S. Nieswand,⁹ R. Niet,¹⁰ N. Nikitin,³³ T. Nikodem,¹² A. Nogay,⁶⁸ D. P. O'Hanlon,⁵⁰ A. Oblakowska-Mucha,²⁸ V. Obraztsov,³⁷ S. Ogilvy,¹⁹ R. Oldeman,^{16,f} C. J. G. Onderwater,⁷¹ A. Ossowska,²⁷ J. M. Otalora Goicochea,² P. Owen,⁴² A. Oyanguren,⁷⁰ P. R. Pais,⁴¹ A. Palano,^{14,l} M. Palutan,^{19,40} A. Papanestis,⁵¹ M. Pappagallo,^{14,l} L. L. Pappalardo,^{17,a} C. Pappenheimer,⁵⁹ W. Parker,⁶⁰ C. Parkes,⁵⁶ G. Passaleva,¹⁸ A. Pastore,^{14,l} M. Patel,⁵⁵ C. Patrignani,^{15,h} A. Pearce,⁴⁰ A. Pellegrino,⁴³ G. Penso,²⁶ M. Pepe Altarelli,⁴⁰ S. Perazzini,⁴⁰ P. Perret,⁵ L. Pescatore,⁴¹ K. Petridis,⁴⁸ A. Petrolini,^{20,j} A. Petrov,⁶⁸ M. Petruzzio,^{22,n} E. Picatoste Olloqui,³⁸ B. Pietrzyk,⁴ M. Pikies,²⁷ D. Pinci,²⁶ A. Pistone,^{20,j} A. Piucci,¹² V. Placinta,³⁰ S. Playfer,⁵² M. Plo Casasus,³⁹ F. Polci,⁸ M. Poli Lener,¹⁹ A. Poluektov,^{50,36} I. Polyakov,⁶¹ E. Polcarpo,² G. J. Pomery,⁴⁸ S. Ponce,⁴⁰ A. Popov,³⁷ D. Popov,^{11,40} S. Poslavskii,³⁷ C. Potterat,² E. Price,⁴⁸ J. Prisciandaro,³⁹ C. Prouve,⁴⁸ V. Pugatch,⁴⁶ A. Puig Navarro,⁴² H. Pullen,⁵⁷ G. Punzi,^{24,t} W. Qian,⁵⁰ R. Quagliani,^{7,48} B. Quintana,⁵ B. Rachwal,²⁸ J. H. Rademacker,⁴⁸ M. Rama,²⁴ M. Ramos Pernas,³⁹ M. S. Rangel,² I. Raniuk,^{45,†} F. Ratnikov,³⁵ G. Raven,⁴⁴ M. Ravonel Salzgeber,⁴⁰ M. Reboud,⁴ F. Redi,⁵⁵ S. Reichert,¹⁰ A. C. dos Reis,¹ C. Remon Alepuz,⁷⁰ V. Renaudin,⁷ S. Ricciardi,⁵¹ S. Richards,⁴⁸ M. Rihl,⁴⁰ K. Rinnert,⁵⁴ V. Rives Molina,³⁸ P. Robbe,⁷ A. Robert,⁸ A. B. Rodrigues,¹ E. Rodrigues,⁵⁹ J. A. Rodriguez Lopez,⁶⁶ P. Rodriguez Perez,^{56,†} A. Rogozhnikov,³⁵

S. Roiser,⁴⁰ A. Rollings,⁵⁷ V. Romanovskiy,³⁷ A. Romero Vidal,³⁹ J. W. Ronayne,¹³ M. Rotondo,¹⁹ M. S. Rudolph,⁶¹ T. Ruf,⁴⁰ P. Ruiz Valls,⁷⁰ J. Ruiz Vidal,⁷⁰ J. J. Saborido Silva,³⁹ E. Sadykhov,³² N. Sagidova,³¹ B. Saitta,^{16,f} V. Salustino Guimaraes,¹ D. Sanchez Gonzalo,³⁸ C. Sanchez Mayordomo,⁷⁰ B. Sanmartin Sedes,³⁹ R. Santacesaria,²⁶ C. Santamarina Rios,³⁹ M. Santimaria,¹⁹ E. Santovetti,^{25,i} G. Sarpis,⁵⁶ A. Sarti,²⁶ C. Satriano,^{26,u} A. Satta,²⁵ D. M. Saunders,⁴⁸ D. Savrina,^{32,33} S. Schael,⁹ M. Schellenberg,¹⁰ M. Schiller,⁵³ H. Schindler,⁴⁰ M. Schlupp,¹⁰ M. Schmelling,¹¹ T. Schmelzer,¹⁰ B. Schmidt,⁴⁰ O. Schneider,⁴¹ A. Schopper,⁴⁰ H. F. Schreiner,⁵⁹ K. Schubert,¹⁰ M. Schubiger,⁴¹ M.-H. Schune,⁷ R. Schwemmer,⁴⁰ B. Sciascia,¹⁹ A. Sciubba,^{26,v} A. Semennikov,³² A. Sergi,⁴⁷ N. Serra,⁴² J. Serrano,⁶ L. Sestini,²³ P. Seyfert,⁴⁰ M. Shapkin,³⁷ I. Shapoval,⁴⁵ Y. Shcheglov,³¹ T. Shears,⁵⁴ L. Shekhtman,^{36,e} V. Shevchenko,⁶⁸ B. G. Siddi,^{17,40} R. Silva Coutinho,⁴² L. Silva de Oliveira,² G. Simi,^{23,q} S. Simone,^{14,l} M. Sirendi,⁴⁹ N. Skidmore,⁴⁸ T. Skwarnicki,⁶¹ E. Smith,⁵⁵ I. T. Smith,⁵² J. Smith,⁴⁹ M. Smith,⁵⁵ I. Soares Lavra,¹ M. D. Sokoloff,⁵⁹ F. J. P. Soler,⁵³ B. Souza De Paula,² B. Spaan,¹⁰ P. Spradlin,⁵³ S. Sridharan,⁴⁰ F. Stagni,⁴⁰ M. Stahl,¹² S. Stahl,⁴⁰ P. Steffko,⁴¹ S. Stefkova,⁵⁵ O. Steinkamp,⁴² S. Stemmler,¹² O. Stenyakin,³⁷ M. Stepanova,³¹ H. Stevens,¹⁰ S. Stone,⁶¹ B. Storaci,⁴² S. Stracka,^{24,t} M. E. Stramaglia,⁴¹ M. Straticiu,³⁰ U. Straumann,⁴² L. Sun,⁶⁴ W. Sutcliffe,⁵⁵ K. Swientek,²⁸ V. Syropoulos,⁴⁴ M. Szczekowski,²⁹ T. Szumlak,²⁸ M. Szymanski,⁶³ S. T'Jampens,⁴ A. Tayduganov,⁶ T. Tekampe,¹⁰ G. Tellarini,^{17,a} F. Teubert,⁴⁰ E. Thomas,⁴⁰ J. van Tilburg,⁴³ M. J. Tilley,⁵⁵ V. Tisserand,⁴ M. Tobin,⁴¹ S. Tolk,⁴⁹ L. Tomassetti,^{17,a} D. Tonelli,²⁴ F. Toriello,⁶¹ R. Tourinho Jadallah Aoude,¹ E. Tournefier,⁴ M. Traill,⁵³ M. T. Tran,⁴¹ M. Tresch,⁴² A. Trisovic,⁴⁰ A. Tsaregorodtsev,⁶ P. Tsopelas,⁴³ A. Tully,⁴⁹ N. Tuning,⁴³ A. Ukleja,²⁹ A. Usachov,⁷ A. Ustyuzhanin,³⁵ U. Uwer,¹² C. Vacca,^{16,f} A. Vagner,⁶⁹ V. Vagnoni,^{15,40} A. Valassi,⁴⁰ S. Valat,⁴⁰ G. Valenti,¹⁵ R. Vazquez Gomez,¹⁹ P. Vazquez Regueiro,³⁹ S. Vecchi,¹⁷ M. van Veghel,⁴³ J. J. Velthuis,⁴⁸ M. Veltri,^{18,w} G. Veneziano,⁵⁷ A. Venkateswaran,⁶¹ T. A. Verlage,⁹ M. Vernet,⁵ M. Vesterinen,⁵⁷ J. V. Viana Barbosa,⁴⁰ B. Viaud,⁷ D. Vieira,⁶³ M. Vieites Diaz,³⁹ H. Viemann,⁶⁷ X. Vilasis-Cardona,^{38,g} M. Vitti,⁴⁹ V. Volkov,³³ A. Vollhardt,⁴² B. Voneki,⁴⁰ A. Vorobyev,³¹ V. Vorobyev,^{36,e} C. Voß,⁹ J. A. de Vries,⁴³ C. Vázquez Sierra,³⁹ R. Waldi,⁶⁷ C. Wallace,⁵⁰ R. Wallace,¹³ J. Walsh,²⁴ J. Wang,⁶¹ D. R. Ward,⁴⁹ H. M. Wark,⁵⁴ N. K. Watson,⁴⁷ D. Websdale,⁵⁵ A. Weiden,⁴² M. Whitehead,⁴⁰ J. Wicht,⁵⁰ G. Wilkinson,^{57,40} M. Wilkinson,⁶¹ M. Williams,⁵⁶ M. P. Williams,⁴⁷ M. Williams,⁵⁸ T. Williams,⁴⁷ F. F. Wilson,⁵¹ J. Wimberley,⁶⁰ M. A. Winn,⁷ J. Wishahi,¹⁰ W. Wislicki,²⁹ M. Witek,²⁷ G. Wormser,⁷ S. A. Wotton,⁴⁹ K. Wraight,⁵³ K. Wyllie,⁴⁰ Y. Xie,⁶⁵ Z. Xu,⁴ Z. Yang,³ Z. Yang,⁶⁰ Y. Yao,⁶¹ H. Yin,⁶⁵ J. Yu,⁶⁵ X. Yuan,⁶¹ O. Yushchenko,³⁷ K. A. Zarebski,⁴⁷ M. Zavertyaev,^{11,b} L. Zhang,³ Y. Zhang,⁷ A. Zhelezov,¹² Y. Zheng,⁶³ X. Zhu,³ V. Zhukov,³³ J. B. Zonneveld,⁵² and S. Zucchelli¹⁵

(LHCb Collaboration)

¹Centro Brasileiro de Pesquisas Físicas (CBPF), Rio de Janeiro, Brazil²Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil³Center for High Energy Physics, Tsinghua University, Beijing, China⁴LAPP, Université Savoie Mont-Blanc, CNRS/IN2P3, Annecy-Le-Vieux, France⁵Clermont Université, Université Blaise Pascal, CNRS/IN2P3, LPC, Clermont-Ferrand, France⁶CPPM, Aix-Marseille Université, CNRS/IN2P3, Marseille, France⁷LAL, Université Paris-Sud, CNRS/IN2P3, Orsay, France⁸LPNHE, Université Pierre et Marie Curie, Université Paris Diderot, CNRS/IN2P3, Paris, France⁹I. Physikalisches Institut, RWTH Aachen University, Aachen, Germany¹⁰Fakultät Physik, Technische Universität Dortmund, Dortmund, Germany¹¹Max-Planck-Institut für Kernphysik (MPIK), Heidelberg, Germany¹²Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany¹³School of Physics, University College Dublin, Dublin, Ireland¹⁴Sezione INFN di Bari, Bari, Italy¹⁵Sezione INFN di Bologna, Bologna, Italy¹⁶Sezione INFN di Cagliari, Cagliari, Italy¹⁷Università e INFN, Ferrara, Ferrara, Italy¹⁸Sezione INFN di Firenze, Firenze, Italy¹⁹Laboratori Nazionali dell'INFN di Frascati, Frascati, Italy²⁰Sezione INFN di Genova, Genova, Italy²¹Università e INFN, Milano-Bicocca, Milano, Italy²²Sezione di Milano, Milano, Italy

- ²³Sezione INFN di Padova, Padova, Italy
²⁴Sezione INFN di Pisa, Pisa, Italy
²⁵Sezione INFN di Roma Tor Vergata, Roma, Italy
²⁶Sezione INFN di Roma La Sapienza, Roma, Italy
²⁷Henryk Niewodniczanski Institute of Nuclear Physics Polish Academy of Sciences, Kraków, Poland
²⁸AGH - University of Science and Technology, Faculty of Physics and Applied Computer Science, Kraków, Poland
²⁹National Center for Nuclear Research (NCBJ), Warsaw, Poland
³⁰Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest-Magurele, Romania
³¹Petersburg Nuclear Physics Institute (PNPI), Gatchina, Russia
³²Institute of Theoretical and Experimental Physics (ITEP), Moscow, Russia
³³Institute of Nuclear Physics, Moscow State University (SINP MSU), Moscow, Russia
³⁴Institute for Nuclear Research of the Russian Academy of Sciences (INR RAN), Moscow, Russia
³⁵Yandex School of Data Analysis, Moscow, Russia
³⁶Budker Institute of Nuclear Physics (SB RAS), Novosibirsk, Russia
³⁷Institute for High Energy Physics (IHEP), Protvino, Russia
³⁸ICCUB, Universitat de Barcelona, Barcelona, Spain
³⁹Universidad de Santiago de Compostela, Santiago de Compostela, Spain
⁴⁰European Organization for Nuclear Research (CERN), Geneva, Switzerland
⁴¹Institute of Physics, Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland
⁴²Physik-Institut, Universität Zürich, Zürich, Switzerland
⁴³Nikhef National Institute for Subatomic Physics, Amsterdam, The Netherlands
⁴⁴Nikhef National Institute for Subatomic Physics and VU University Amsterdam, Amsterdam, The Netherlands
⁴⁵NSC Kharkiv Institute of Physics and Technology (NSC KIPT), Kharkiv, Ukraine
⁴⁶Institute for Nuclear Research of the National Academy of Sciences (KINR), Kyiv, Ukraine
⁴⁷University of Birmingham, Birmingham, United Kingdom
⁴⁸H.H. Wills Physics Laboratory, University of Bristol, Bristol, United Kingdom
⁴⁹Cavendish Laboratory, University of Cambridge, Cambridge, United Kingdom
⁵⁰Department of Physics, University of Warwick, Coventry, United Kingdom
⁵¹STFC Rutherford Appleton Laboratory, Didcot, United Kingdom
⁵²School of Physics and Astronomy, University of Edinburgh, Edinburgh, United Kingdom
⁵³School of Physics and Astronomy, University of Glasgow, Glasgow, United Kingdom
⁵⁴Oliver Lodge Laboratory, University of Liverpool, Liverpool, United Kingdom
⁵⁵Imperial College London, London, United Kingdom
⁵⁶School of Physics and Astronomy, University of Manchester, Manchester, United Kingdom
⁵⁷Department of Physics, University of Oxford, Oxford, United Kingdom
⁵⁸Massachusetts Institute of Technology, Cambridge, Massachusetts, USA
⁵⁹University of Cincinnati, Cincinnati, Ohio, USA
⁶⁰University of Maryland, College Park, Maryland, USA
⁶¹Syracuse University, Syracuse, New York, USA
⁶²Pontifícia Universidade Católica do Rio de Janeiro (PUC-Rio), Rio de Janeiro, Brazil
 (associated with Institution Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil)
⁶³University of Chinese Academy of Sciences, Beijing, China
 (associated with Institution Center for High Energy Physics, Tsinghua University, Beijing, China)
⁶⁴School of Physics and Technology, Wuhan University, Wuhan, China
 (associated with Institution Center for High Energy Physics, Tsinghua University, Beijing, China)
⁶⁵Institute of Particle Physics, Central China Normal University, Wuhan, Hubei, China
 (associated with Institution Center for High Energy Physics, Tsinghua University, Beijing, China)
⁶⁶Departamento de Física, Universidad Nacional de Colombia, Bogota, Colombia
 (associated with Institution LPNHE, Université Pierre et Marie Curie, Université Paris Diderot, CNRS/IN2P3, Paris, France)
⁶⁷Institut für Physik, Universität Rostock, Rostock, Germany
 (associated with Institution Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany)
⁶⁸National Research Centre Kurchatov Institute, Moscow, Russia
 (associated with Institution Institute of Theoretical and Experimental Physics (ITEP), Moscow, Russia)
⁶⁹National Research Tomsk Polytechnic University, Tomsk, Russia
 (associated with Institution Institute of Theoretical and Experimental Physics (ITEP), Moscow, Russia)
⁷⁰Instituto de Física Corpuscular, Centro Mixto Universidad de Valencia - CSIC, Valencia, Spain
 (associated with Institution ICCUB, Universitat de Barcelona, Barcelona, Spain)
⁷¹Van Swinderen Institute, University of Groningen, Groningen, The Netherlands
 (associated with Institution Nikhef National Institute for Subatomic Physics, Amsterdam, The Netherlands)

[†]Deceased.

^aAlso at Università di Ferrara, Ferrara, Italy.

^bAlso at P.N. Lebedev Physical Institute, Russian Academy of Science (LPI RAS), Moscow, Russia.

^cAlso at Università di Milano Bicocca, Milano, Italy.

^dAlso at Università di Modena e Reggio Emilia, Modena, Italy.

^eAlso at Novosibirsk State University, Novosibirsk, Russia.

^fAlso at Università di Cagliari, Cagliari, Italy.

^gAlso at LIFAELS, La Salle, Universitat Ramon Llull, Barcelona, Spain.

^hAlso at Università di Bologna, Bologna, Italy.

ⁱAlso at Università di Roma Tor Vergata, Roma, Italy.

^jAlso at Università di Genova, Genova, Italy.

^kAlso at Scuola Normale Superiore, Pisa, Italy.

^lAlso at Università di Bari, Bari, Italy.

^mAlso at Laboratoire Leprince-Ringuet, Palaiseau, France.

ⁿAlso at Università degli Studi di Milano, Milano, Italy.

^oAlso at Universidade Federal do Triângulo Mineiro (UFTM), Uberaba-MG, Brazil.

^pAlso at AGH - University of Science and Technology, Faculty of Computer Science, Electronics and Telecommunications, Kraków, Poland.

^qAlso at Università di Padova, Padova, Italy.

^rAlso at Iligan Institute of Technology (IIT), Iligan, Philippines.

^sAlso at Hanoi University of Science, Hanoi, Viet Nam.

^tAlso at Università di Pisa, Pisa, Italy.

^uAlso at Università della Basilicata, Potenza, Italy.

^vAlso at Università di Roma La Sapienza, Roma, Italy.

^wAlso at Università di Urbino, Urbino, Italy.